

Ciprofloxacin-Releasing Bioabsorbable Polymer is Superior to Titanium in Preventing *Staphylococcus Epidermidis* Attachment and Biofilm Formation in Vitro.

Sanna-Mari Niemelä¹, Irma Ikäheimo², Markku Koskela², Minna Veiranto³, Esa Suokas⁴, Pertti Törmälä³, Timo Waris¹, Nureddin Ashammakhi^{1,3}, Hannu Syrjälä⁵

1. Department of Surgery, Oulu University Hospital, Oulu, Finland.
2. Clinical Microbiology Laboratory of Oulu University Hospital, Oulu, Finland.
3. Institute of Biomaterials, Tampere University of Technology, Tampere, Finland.
4. Linvatec Biomaterials Ltd., Tampere, Finland.
5. Department of Infection Control, Oulu University Hospital, Oulu, Finland.

Background: Antibiotic coating systems have been successfully used to prevent bacterial attachment and biofilm formation. Our purpose was to evaluate whether bioabsorbable polylactide-co-glycolide (PLGA) 80/20 on its own, and PLGA together with ciprofloxacin (PLGA+AB) have any advantages over titanium in preventing *Staphylococcus epidermidis* attachment and biofilm formation in vitro.

Materials and methods: Cylindrical specimens of titanium, PLGA and PLGA+AB in triplicate were examined for *S. epidermidis* ATCC 35989 attachment and biofilm formation after incubation with a bacterial suspension of ca. 10⁵ cfu/ml for 1, 3, 7, 14 and 21 days, using scanning electron microscopy. Growth inhibition properties of PLGA and PLGA + AB cylinders were tested on agar plates.

Results: On days 1, 3 and 21, no bacterial attachment was seen in 19.5%, 9.2% and 41.4% of the titanium specimens, in 18.4%, 28.7% and 34.5% of the PLGA specimens and in 57.5%, 62.1% and 57.5% of the PLGA + AB specimens, respectively. During the whole study period no biofilm was observed on 74%–93% of the titanium specimens, 58%–78% of the PLGA specimens and 93%–100% of the PLGA+AB specimens. PLGA + AB showed clear bacterial growth inhibition on agar plates while PLGA and titanium did not show any inhibition.

Conclusions: PLGA + AB bioabsorbable material was superior to titanium in preventing bacterial attachment and biofilm formation and may have clinical applicability, for example, in prevention of infection in trauma surgery or in the treatment of chronic osteomyelitis.

Keywords: antibiotic, bioabsorbable, biofilm, ciprofloxacin, polylactide-co-glycolide

